


# One-Parameter Deformations of Bowen-Series Functions Associated To Cocompact Fuchsian Triangle Groups

---

Ayşe Yıltekin Karataş  
Oregon State University  
Numeration Conference, May 2023



# Motivation

---

# Motivation

- Relationship between the Regular Continued Fraction map  $T$  and the action of  $SL_2(\mathbb{Z})$  on  $\mathbb{R}$ .
- The RCF map  $T$  is expansive, Markov, transitive and satisfy Rényi's condition.
- Gauss measure:  $T$ -invariant and equivalent to Lebesgue
- Nakada's  $\alpha$ -continued fraction maps  $T_\alpha$  give a one-parameter deformation of  $T$ .

## Motivation

Let  $\Gamma$  be a finitely generated discrete subgroup of  $SL_2(\mathbb{R})$  acting on  $\mathbb{R}$  with dense orbits; i.e. finitely generated Fuchsian group of the first kind.

RCF	B-S
$SL_2(\mathbb{Z})$ acting on $\mathbb{R}$	$\Gamma$ acting on $\mathbb{D}$
$T : (0,1) \rightarrow (0,1)$	$f: \mathbb{S}^1 \rightarrow \mathbb{S}^1$
Expansivity, Markov	✓
Transitive, Rényi's condition	✓
Gauss measure	✓
$T_\alpha$	Analog ?

Rufus Bowen and Caroline Series, *Markov Maps Associated with Fuchsian Groups*, 1979

- Katok and Ugarcovici (2017) studied B-S functions associated to cocompact torsion-free Fuchsian groups and defined a multi-parameter deformation family.
- Los (2009) defined Bowen-Series like maps for cocompact surface groups considering the geometric presentation of the group. This study excludes triangle groups.

# Main Results

The following is joint work with my Ph.D. advisor, Thomas A. Schmidt.

- Correction to Bowen & Series (1979).

- (Q1) Can we define a family of expansive functions via an  $\alpha$ -deformation of  $f$  in the case of cocompact Fuchsian triangle groups.?
- (Q2) If so, is there an ergodic invariant measure for each function in the family?

# The Bowen-Series Fundamental Domain

---

# Cocompact Fuchsian Triangle Groups

- A **cocompact Fuchsian triangle group**  $\Gamma$  is a group with signature  $(0; m_1, m_2, m_3)$ , where  $m_i \in \mathbb{Z}^+$  and  $\frac{1}{m_1} + \frac{1}{m_2} + \frac{1}{m_3} < 1$ .

## Example

$(6,6,3)$  with the presentation

$$\{A, B \mid A^6 = B^6 = (AB)^3 = I\}.$$

- The elements in  $\Gamma$ , as Möbius transformations, act on the unit disc.



# Cocompact Fuchsian Triangle Groups

- A **cocompact Fuchsian triangle group**  $\Gamma$  is a group with signature  $(0; m_1, m_2, m_3)$ , where  $m_i \in \mathbb{Z}^+$  and  $\frac{1}{m_1} + \frac{1}{m_2} + \frac{1}{m_3} < 1$ .

## Example


$(6,6,3)$  with the presentation

$$\{A, B \mid A^6 = B^6 = (AB)^3 = I\}.$$

- The elements in  $\Gamma$ , as Möbius transformations, act on the unit disc.
- The elements in  $\Gamma$  are classified as hyperbolic and elliptic. A **hyperbolic** element has **two fixed points on  $\mathbb{S}^1$**  and an **elliptic** element has a **single fixed point on the interior of the unit disc**.


## Constructing a Fundamental Domain

A closed region  $\mathcal{F} \subset \mathbb{D}$  is called **fundamental domain** for  $\Gamma$  if it tessellates  $\mathbb{D}$  under the action of  $\Gamma$ .



**Figure 1:** A fundamental domain  $\mathcal{F}$  for the triangle group  $(6,6,3)$ .

# Constructing a Fundamental Domain



**Figure 2:** A fundamental domain  $\mathcal{F}$  for the triangle group  $(6,6,3)$

# Constructing a Fundamental Domain

A fundamental domain  $\mathcal{F}$  has **extension property** if for all  $s \in S$ ,


$$g(s) \cap \bigcup_{T \in \Gamma} T(\mathcal{F}^\circ) = \emptyset,$$

where  $g(s)$  represents the geodesic containing the side  $s$ .

## Constructing a Fundamental Domain

We consider the set  $N = \cup_{i=1}^4 N_i$  in  $\mathbb{D}$ , where  $N_i$  is defined as


$N_i := \{g \text{ geodesics} \mid v_i \in g \text{ and } \exists T \in \Gamma, T(s_j) \subset g \text{ for some } j\}$ .



**Figure 3:**  $N$  for the group of signature  $(6,6,3)$ .

## Correction to Bowen & Series (1979)

For  $\mathcal{F}$  to satisfy the **extension property**, no geodesic of any  $N_i$  meets  $\mathcal{F}^\circ$ .



**Figure 4:**  $N$  for the group of signature (3,5,6).

## Correction to Bowen & Series (1979)

### Theorem (A.Y.K. & Schmidt, 2023)

Suppose that  $(m_1, m_2, m_3)$  is the signature of a cocompact hyperbolic Fuchsian triangle group. If more than one  $m_i$  is odd, then no convex fundamental domain for the signature has the extension property. Otherwise, the Bowen-Series fundamental domain for this signature does have the extension property.

In what follows, we suppose that the Bowen-Series fundamental domain for  $\Gamma$  satisfies extension property.

# One-Parameter Deformation of B-S Functions


---



# The Bowen-Series function

The **Bowen - Series function**  $f$  is defined as

$$f(x) := T_i(x) \text{ on } x \in [a_i, a_{i+1}).$$




**Figure 5:** The Bowen-Series function  $f$  for the group  $(6,6,3)$ . Graphed as  $y = \arg(f(x))$  for  $x \in [0, 2\pi)$ .

## A Function Family


We call  $\mathcal{O}_i = [a_i, b_i)$  as the **overlap interval** and define a function family depending on the parameter  $\alpha$  as

$$f_\alpha(x) := \begin{cases} f(x), & x \in \mathbb{S}^1 \setminus \mathcal{D} \\ T_{i-1}(x), & x \in \mathcal{D}, \end{cases}$$

where  $\mathcal{D} = [a_i, \alpha)$  is the **differing interval**.



# A Function Family



**Figure 6:** Comparison of the plots of  $f$  and  $f_\alpha$

## Back to Questions

- (Q1) Can we define a family of expansive functions via an  $\alpha$ -deformation of  $f$  in the case of cocompact Fuchsian triangle groups.? Yes
- (Q2) If so, is there an ergodic invariant measure for each function in the family?

## Adler-Flatto's 'Folklore Theorem'

Let  $X$  be an interval or a circle and  $\mathcal{P} = \{I_k\}$  be a finite partition of  $X$ . Let  $g : X \rightarrow X$  satisfy the following:

- (1) Piecewise strict monotonicity,
- (2) Piecewise smoothness,
- (3)  $g(\overline{I_k})$  is equal to the union of some  $\overline{I_\ell}$ 's.
- (4) There exists an integer  $p$  such that  $g^p(\overline{I_k}) = X$  for all  $k$ .

The conditions (1)-(3) implies that  $g$  is a **Markov** map. Condition (4) is called aperiodicity or **transitivity** condition.

## Adler-Flatto's 'Folklore Theorem'


Let  $X$  be an interval or a circle and  $\mathcal{P} = \{I_k\}$  be a finite partition of  $X$ . Let  $g : X \rightarrow X$  satisfy

- (1) Piecewise strict monotonicity,
- (2) Piecewise smoothness,
- (3)  $g(\overline{I_k})$  is equal to the union of some  $\overline{I_\ell}$ 's.
- (4) There exists an integer  $p$  such that  $g^p(\overline{I_k}) = X$  for all  $k$ .

### Theorem

Assume that (1)-(4) hold and  $g$  is eventually expansive. Then  $g$  has an ergodic invariant measure equivalent to Lebesgue measure.


## Partition $\mathcal{P}$ for $f$



**Figure 7:** The set  $E$  of the end points of the geodesics in  $N$  form the partition intervals  $\{I_k\}$ .

## Markov Property for $f$

Bowen & Series show that  $f$  is a Markov function with respect to  $\mathcal{P}$ . To show (3):  $f(\overline{I_k})$  is equal to the union of some  $\overline{I_\ell}$ 's, it is enough to show that  $E$  is invariant under  $f$ .




**Figure 8:** The Bowen-Series function  $f$  for the triangle group  $(6,6,3)$ .



## Partition $\mathcal{P}_\alpha$ for $f_\alpha$

For  $f_\alpha$ , we define  $\mathcal{P}_\alpha$  using

$$E_\alpha = E \cup \{f_\alpha^k(\alpha)\}_{k \geq 0} \cup \{f_\alpha^k(T_{i-1}(\alpha))\}_{k \geq 0}.$$



**Figure 9:** The plot of  $f_\alpha$ . Gridlines correspond to the points in  $E_\alpha$ .

## Markov Property for $f_\alpha$

$E_\alpha$  is  $f_\alpha$ -invariant. Thus,  $f_\alpha$  is Markov if  $E_\alpha$  is finite.

Since  $E$  is finite, we only need that the orbits  $\{f_\alpha^k(\alpha)\}_{k \geq 0}$  and  $\{f_\alpha^k(T_{i-1}(\alpha))\}_{k \geq 0}$  are finite.

### Theorem (A.Y.K. & Schmidt, 2023)

The function  $f_\alpha$  is Markov if and only if  $\alpha$  is a hyperbolic fixed point of  $\Gamma$ .

### Proof

( $\Rightarrow$ ) If  $f_\alpha$  is Markov, then  $E_\alpha$  is finite, then  $f_\alpha$ -orbit of  $\alpha$  is finite, which implies that  $\alpha$  is  $f_\alpha$ -preperiodic; i.e. there exists  $m, n \geq 0$  such that  $f_\alpha^m(\alpha) = f_\alpha^n(\alpha)$ . Hence,  $\alpha$  is a hyperbolic fixed point.

## Markov Property for $f_\alpha$

$E_\alpha$  is  $f_\alpha$ -invariant. Thus,  $f_\alpha$  is Markov if  $E_\alpha$  is finite.

Since  $E$  is finite, we only need that the orbits  $\{f_\alpha^k(\alpha)\}_{k \geq 0}$  and  $\{f_\alpha^k(T_{i-1}(\alpha))\}_{k \geq 0}$  are finite.

### Theorem (A.Y.K. & Schmidt, 2023)

The function  $f_\alpha$  is Markov if and only if  $\alpha$  is a hyperbolic fixed point of  $\Gamma$ .

### Proof

( $\Rightarrow$ ) If  $f_\alpha$  is Markov, then  $E_\alpha$  is finite, then  $f_\alpha$ -orbit of  $\alpha$  is finite, which implies that  $\alpha$  is  $f_\alpha$ -preperiodic; i.e. there exists  $m, n \geq 0$  such that  $f_\alpha^m(\alpha) = f_\alpha^n(\alpha)$ . Hence,  $\alpha$  is a hyperbolic fixed point.

( $\Leftarrow$ ) Suppose  $\alpha$  is a hyperbolic fixed point with infinite  $f_\alpha$ -orbit.

# Markov Property for $f_\alpha$

## Lemma (A.Y.K. & Schmidt, 2023)

Fix  $\alpha \in \mathcal{O}$ . Suppose that  $x \in \mathbb{S}^1$  has infinite  $f_\alpha$ -orbit. Then there are infinitely many values of  $j$  such that the  $f$ -orbit of  $x$  contains either (1)  $f_\alpha^j(x)$ , (2)  $f \circ f_\alpha^j(x)$ , or (3)  $f^2 \circ f_\alpha^j(x)$ .

## Proof (Cont.):

# Markov Property for $f_\alpha$

## Lemma (A.Y.K. & Schmidt, 2023)

Fix  $\alpha \in \mathcal{O}$ . Suppose that  $x \in \mathbb{S}^1$  has infinite  $f_\alpha$ -orbit. Then there are infinitely many values of  $j$  such that the  $f$ -orbit of  $x$  contains either (1)  $f_\alpha^j(x)$ , (2)  $f \circ f_\alpha^j(x)$ , or (3)  $f^2 \circ f_\alpha^j(x)$ .

## Proof (Cont.):

- By the Pigeonhole Principle, we can assume one of the three cases occurs for infinitely many values of  $j$ .

# Markov Property for $f_\alpha$

## Lemma (A.Y.K. & Schmidt, 2023)

Fix  $\alpha \in \mathcal{O}$ . Suppose that  $x \in \mathbb{S}^1$  has infinite  $f_\alpha$ -orbit. Then there are infinitely many values of  $j$  such that the  $f$ -orbit of  $x$  contains either (1)  $f_\alpha^j(x)$ , (2)  $f \circ f_\alpha^j(x)$ , or (3)  $f^2 \circ f_\alpha^j(x)$ .

## Proof (Cont.):

- By the Pigeonhole Principle, we can assume one of the three cases occurs for infinitely many values of  $j$ .
- One easily shows  $f$ -orbit of  $\alpha$  is finite.
- Since  $f$  is a finite-to-one function and the  $f$ -orbit of  $\alpha$  is finite, there are only finitely many preimages under  $f$  or  $f^2$  of this finite set.
- There are some  $m \neq n$  such that  $f_\alpha^m(\alpha) = f_\alpha^n(\alpha)$ , and so  $f_\alpha$ -orbit of  $\alpha$  is finite.



## Theorem (A.Y.K. & Schmidt, 2023)

Fix  $\alpha \in \mathcal{O}_i$ . Let  $n_i = |N_i|$ . The function  $f_\alpha$  is **surjective** if and only if the following conditions hold:

- $n_i > 2$ ,
- $n_i = 2$  and  $n_{i+2} > 2$ ,
- $\alpha$  belongs to the closure of the set of points  $x \in \mathcal{O}_i$  such that  $f^{n_i}(x) = f^{n_i-1}(T_{i-1}x)$ .

Moreover, if  $f_\alpha$  is Markov, then  $f_\alpha$  has ergodic invariant measure equivalent to Lebesgue measure if and only if  $f_\alpha$  is surjective.

# Transitivity



**Figure 10:** Plot of the function  $f_\alpha$  for the signature  $(4, 4, 3)$ . This function is not transitive; it is not even a surjective function!



Thank you for your attention!



**T.A. Schmidt and A. Yıltekin-Karataş,**  
*Continuous Deformation of The Bowen-Series Map Associated to a Cocompact Triangle Group,*  
<https://arxiv.org/abs/2305.04892>, (2023).



**R. Bowen and C. Series,**  
*Markov Maps Associated with Fuchsian Groups,*  
Publications mathématiques de l'I.H.É.S., 50 (1979), p. 153 - 170.



**S. Katok and I. Ugarcovici,**  
*Structure of attractors for boundary maps associated to Fuchsian groups,* Geometriae Dedicata, 191 (2017), p. 171 - 198.



**R. Adler and L. Flatto,**

*Geodesic flows, interval maps and symbolic dynamics,*  
Bulletin of the American Mathematical Society, 25, (1991), no. 2,  
229-334.



**S. Katok,**

*Fuchsian Groups,*  
University of Chicago Press, 1992.



**J. Los,**

*Volume entropy for surface groups via Bowen-Series-like maps,*  
J. Topol. 7 (2014), no. 1, 120 - 154.



<http://www.malinc.se/m/ImageTiling.php>